

Electromagnetic Plane Waves (Cont'd)

Wave Packets

By superposing plane waves with different frequencies, one can obtain wave packets that last for a finite time and have finite spatial extension. Ignoring polarization of the wave, a wave packet can be written in the following form (assuming one-dimensional space):

$$U(x, t) = \int A(k) e^{i(kx - \omega t)} dk$$

For non-dispersive media, $k = \frac{\omega}{c} n$, where n is constant. Then:

$$U(x, t) = \int A(k) e^{ik(x - \frac{c}{n}t)} dk = U(x - vt, 0) \quad (v = \frac{c}{n})$$

In this case, the wave packet is just translated in space without any change in its shape.

For dispersive media, $n = n(\omega)$ and $k(\omega) = \frac{\omega}{c} n(\omega)$. Assuming that $A(k)$ is significant over a small range of k around k_0 , we have:

$$\omega(k) = \omega(k_0) + \omega'(k_0)(k - k_0) + \frac{\omega''(k_0)}{2}(k - k_0)^2 + \dots$$

Then:

$$U(x, t) = \int A(k) e^{i[kx - \omega(k_0)t - \omega'(k_0)(k - k_0)t + \dots]} dk \Rightarrow U(x, t) = e^{i[k_0 x - \omega(k_0)t]} \int A(k) e^{i(kx - v_g t)} dk$$

Here $v_g \equiv \omega'(k_0)$ is called the "group velocity". Therefore:

$$U(x, t) = e^{i(k_0 v_g x - \omega_0 t)} U(x - v_g t, 0) \quad (\omega_0 \equiv \omega(k_0))$$

This is just a plane wave modulated by a wave packet that moves at speed v_g without distortion.

Next, let us consider the second-order term $\frac{\omega''(k_0)}{2}(k - k_0)^2$. Then:

$$U(x, t) = e^{i(k_0 v_g x - \omega_0 t)} \int A(k) e^{i(kx - v_g t) - \frac{i}{2}(k - k_0)^2 \omega''(k_0)} dk$$

The last term in the integral leads to spreading of the wave

packet. To see this in more detail, we consider the following example.

Example: Wave packet with a Gaussian profile.

$$A(k) = e^{-\frac{\sigma^2}{2}(k - k_0)^2}$$

Thus:

$$U(\eta, t) = e^{i(k_0 v_g \eta - \omega_0 t)} \int_{-\infty}^{+\infty} e^{-\frac{\sigma^2}{2} (k-k_0)^2 - \frac{i}{2} (k-k_0)^2 \omega''(k_0) t + i(k-k_0)(\eta - v_g t)} dk$$

The integral becomes:

$$\int_{-\infty}^{+\infty} e^{-\left(\frac{\sigma^2}{2} + \frac{i}{2} \omega''(k_0) t\right) (k-k_0)^2 + i(k-k_0)(\eta - v_g t)} dk$$

By completing the square in the exponent, we find:

$$\int_{-\infty}^{+\infty} e^{-\frac{1}{2} (\sigma^2 + i \omega''(k_0) t) (k-k_0)^2 + i(k-k_0)(\eta - v_g t)} dk = \sqrt{\frac{2\pi}{\sigma^2 + i \omega''(k_0) t}}$$

$$e^{i(k-k_0)(\eta - v_g t)} = \sqrt{\frac{2\pi}{\sigma^2 + i \omega''(k_0) t}} \exp\left[-\frac{(\eta - v_g t)^2}{\sigma^2 + i \omega''(k_0) t}\right]$$

$$= \sqrt{\frac{2\pi}{\sigma^2 + i \omega''(k_0) t}} \exp\left[-\frac{(\eta - v_g t)^2}{\sigma^2 + \frac{\omega''(k_0) t}{\sigma^2}}\right] \exp\left[\frac{i}{2} \frac{\omega''(k_0) (\eta - v_g t)^2}{\sigma^2 + \omega''(k_0) t}\right]$$

Hence:

$$|U(\eta, t)| \sim \frac{(2\pi)^{1/2}}{(\sigma^2 + \omega''(k_0) t)^{1/4}} \exp\left[-\frac{(\eta - v_g t)^2}{\sigma^2 + \frac{\omega''(k_0) t}{\sigma^2}}\right]$$

The width of the wave packet at time t is:

$$\sigma(t) = \sqrt{\sigma^2 + \frac{\omega''(k_0) t}{\sigma^2}}$$

At this time the center of the wave packet has traveled a distance $v_g t$. We see that the narrower the wave packet is, i.e., the larger σ is, the faster it spreads. It takes a time $\sim \frac{\sigma^2}{\omega''(k_0)}$ for the wave packet to spread to $\sqrt{2}$ of its original width. The quantity $\omega''(k_0)$ is thus called the "group-velocity dispersion".

One can show that the envelope of $u(x, t)$ with the quadratic dispersion approximation obey a Schrödinger-like equation in the variables t and $X \equiv x - v_g t$.