

## Review of Electrostatics (Cont'd)

### Electrostatics in Dielectrics

When an electric field is applied to a medium, the atomic/molecular constituents have bound charges that get displaced and their centers separate. This is the "polarization" process, which results in formation of electric dipole moments throughout the medium.

Polarization results in volume and surface densities for the bound charge,  $\rho_{\text{bound}}$  and  $\sigma_{\text{bound}}$  respectively, according to:

$$\vec{\nabla} \cdot \vec{P} = -\rho_{\text{bound}}, \quad \vec{P} \cdot \hat{n} = \sigma_{\text{bound}}$$

Here  $\vec{P}$  is the electric dipole moment and  $\hat{n}$  is the unit vector normal to the surface of the dielectric. Since  $\vec{E}$  is related to

the total charge density  $\rho_{\text{tot}} = \rho + \rho_{\text{bound}}$ , we have:

$$\vec{\nabla} \cdot \vec{E} = \frac{1}{\epsilon_0} (\rho - \vec{\nabla} \cdot \vec{P}) \Rightarrow \vec{\nabla} \cdot (\epsilon_0 \vec{E} + \vec{P}) = \rho$$

For linear response of the medium, we have:

$$\vec{P} = \epsilon_0 \chi_e \vec{E} \quad (\chi_e: \text{electric susceptibility})$$

Then:


$$\vec{D} \equiv \epsilon_0 \vec{E} + \vec{P} = \epsilon_0 (1 + \chi_e) \vec{E} = \epsilon \vec{E} \Rightarrow \nabla \cdot \vec{D} = \rho, \quad \nabla \cdot \vec{E} = \frac{\rho}{\epsilon}$$

Here  $\epsilon \equiv \epsilon_0 (1 + \chi_e)$  is electric permittivity of the medium, and

$\frac{\epsilon}{\epsilon_0} = 1 + \chi_e$  is the dielectric constant of the medium.

The boundary conditions at the interface of dielectric media

follow from  $\nabla \cdot \vec{D} = \rho$  and  $\nabla \times \vec{E} = 0$ :

$$\begin{cases} D_{2,n} - D_{1,n} = \sigma \\ E_{2,t} - E_{1,t} = 0 \end{cases} \Rightarrow \begin{cases} \epsilon_2 E_{2,n} - \epsilon_1 E_{1,n} = \sigma \\ E_{2,t} = E_{1,t} \end{cases}$$


Note that  $\epsilon = \epsilon_0$  corresponds to the vacuum and  $\epsilon \rightarrow 0$  describes a conductor.

The electrostatic energy in a dielectric medium follows the

general expression  $U = \frac{1}{2} \int_V \rho \Phi dV$ . After using  $\nabla \cdot \vec{D} = \rho$  and

$\vec{E} = -\vec{\nabla}\Phi$ , it can be shown that:

$$U = \int_V \frac{1}{2} (\vec{D} \cdot \vec{E}) d\tau$$

Thus the electrostatic energy density is given by  $\frac{1}{2} \vec{D} \cdot \vec{E}$

(which is  $\frac{1}{2} \epsilon |\vec{E}|^2 = \frac{|\vec{D}|^2}{2\epsilon}$ ) inside a dielectric.

We now consider examples of boundary value problems involving

dielectric bodies. An important difference from problems with


conducting bodies is that the boundary is not an equipotential

surface in this case. Instead, one has to solve the outside and

inside problems and use the conditions  $E_{in,t} = E_{out,t}$  and

$$\epsilon_{out} E_{out,n} - \epsilon_{in} E_{in,n} = \sigma$$
 to match the solutions at the boundary.

Example: Dielectric sphere in a uniform external electric field.



Inside the sphere,  $\vec{D} = \epsilon \vec{E}$  and  $S_{\text{tot}} = 0$ . Hence,  $\vec{\nabla} \cdot \vec{E} = \frac{1}{\epsilon} \vec{\nabla} \cdot \vec{D} = 0$ ,

and  $\Phi_{\text{in}}$  follows:

$$\Phi_{\text{in}} = \sum_{l=0}^{\infty} A_l r^l P_l(\cos\theta)$$

Here, we have used the fact that  $\Phi_{\text{in}}$  must be finite at the origin. Also, with our choice of z axis, there is no  $\phi$  dependence, which implies that only terms with  $m=0$  are present in the expression (note that  $Y_{l,0}(\theta, \phi) = P_l(\cos\theta)$ ).

Outside the sphere,  $S_{\text{tot}} = 0$ , and therefore:

$$\Phi_{\text{out}} = \sum_{l=0}^{\infty} [B_l r^l + C_l r^{-(l+1)}] P_l(\cos\theta)$$

The asymptotic behavior of  $\Phi_{\text{out}}$  at  $r \rightarrow \infty$  must be that of a uniform field  $E_0$ . Hence,  $B_l = 0$  for  $l \geq 2$  and  $B_1 = -E_0$ .

The individual coefficients that are left will be determined by using the following relations:

$$E_{\text{in},t} = E_{\text{out},t} \quad , \quad \epsilon_0 E_{\text{out},n} \stackrel{!}{=} \epsilon E_{\text{in},n} \quad (\text{since there is no free charge})$$

We note that at  $r=a$  we have:

$$E_t = E_\theta = -\frac{1}{a} \frac{\partial \Phi}{\partial \theta} \Big|_{r=a}, \quad E_n = E_r = -\frac{\partial \Phi}{\partial r} \Big|_{r=a}$$

After using the expressions for  $\Phi_{in}$  and  $\Phi_{out}$  on the previous page, these relations result in:

$$A_1 = \left( \frac{-3}{2 + \frac{\epsilon}{\epsilon_0}} \right) E_0, \quad C_1 = \left( \frac{\frac{\epsilon}{\epsilon_0} - 1}{\frac{\epsilon}{\epsilon_0} + 2} \right) a^3 E_0, \quad A_2 = C_2 = 0 \quad \epsilon \gg 2$$

Therefore:

$$\Phi_{in} = \left( \frac{-3}{2 + \frac{\epsilon}{\epsilon_0}} \right) E_0 r \cos \theta, \quad \Phi_{out} = -E_0 r \cos \theta + \left( \frac{\frac{\epsilon}{\epsilon_0} - 1}{\frac{\epsilon}{\epsilon_0} + 2} \right) E_0 \frac{a^3}{r^2} \cos \theta$$

We thus have a uniform electric field of strength  $\frac{3}{2 + \frac{\epsilon}{\epsilon_0}} E_0$

inside the sphere, resulting in the following polarization:


$$\vec{P} = (\epsilon - \epsilon_0) \vec{E} = 3\epsilon_0 \left( \frac{\frac{\epsilon}{\epsilon_0} - 1}{\frac{\epsilon}{\epsilon_0} + 2} \right) E_0 \hat{z}$$

Note that in the limit  $\epsilon \rightarrow \infty$ , we have  $\vec{E}_{in} = 0$ , as expected for

a conducting sphere. In this case,  $\vec{E}_{out}$  matches the result that

we obtained for a conducting sphere in a uniform electric field.

Example: Point charge  $q$  in front of a dielectric sphere.



The potential outside the sphere follows:

$$\Phi_{out} = \frac{q}{4\pi\epsilon_0 |\vec{x} - \vec{x}_0|} + \sum_{l=0}^{\infty} B_l r^{-(l+1)} P_l(\cos\theta)$$

The first term is due to the charge  $q$  and the second term represents the effect of bound charges on the surface of the sphere. Note that, due to localized nature of the distribution,

$$\Phi_{out} \rightarrow 0 \text{ as } r \rightarrow \infty.$$

The potential inside the sphere is given by:

$$\Phi_{in} = \sum_{l=0}^{\infty} A_l r^l P_l(\cos\theta)$$

Here only positive powers of  $r$  are present so that  $\Phi_{in}$  remains finite at  $r=0$ . We note that  $\Phi_{in}$  is a solution to the Laplace equation as  $\rho_{tot} = 0$  anywhere inside the sphere.

Again, we match the inside and outside solutions at  $r=a$

according to:

$$-\frac{1}{a} \frac{\partial \Phi_{in}}{\partial \theta} \Big|_{r=a} = -\frac{1}{a} \frac{\partial \Phi_{out}}{\partial \theta} \Big|_{r=a}, \quad -\epsilon \frac{\partial \Phi_{in}}{\partial r} \Big|_{r=a} = -\epsilon_0 \frac{\partial \Phi_{out}}{\partial r} \Big|_{r=a}$$

At  $r=a$ , we have:

$$\frac{1}{|\vec{x} - \vec{x}_0|} = \sum_{l=0}^{\infty} \frac{r^l}{d^{l+1}} P_l(\cos \theta)$$

The first condition above (coming from  $E_{in,t} = E_{out,t}$ ) gives:

$$A_l a^l = B_l a^{-(l+1)} + \frac{q a^l}{4\pi \epsilon_0 d^{l+1}}$$

The second condition results in:

$$\epsilon l A_l a^{l-1} = -\epsilon_0 B_l (l+1) a^{-(l+2)} + \frac{q}{4\pi} \frac{l a^{l-1}}{d^{l+1}}$$

Then:

$$A_l = \frac{q l}{4\pi [\epsilon l + \epsilon_0 (l+1)] d^{l+1}}, \quad B_l = -\frac{q a^{2l+1} (\epsilon_0 + \epsilon l)}{4\pi \epsilon_0 [\epsilon l + \epsilon_0 (l+1)] d^{l+1}}$$

Substituting these in the expressions for  $\Phi_{in}$  and  $\Phi_{out}$  allows us to determine the potential everywhere in the space.